INDEX

Acid, mineral, use of, for preserving fodder, 144 sqq.
Africa, East, agricultural grasses in, 341–6.
— Elephant grass in, 345.
— red soils in, 107.
— root-systems of *Coffea arabica* in, 270–84, 285–96.
Africa, South, manuring of sugar-cane in, 379.
— red soils in, 103.
Africa, West, red soils in, 107.
America, Central, red soils in, 104.
America, North, manuring of sugar-cane, 379.
America, South, manuring of sugar-cane, 379.
Ammonium sulphate for sugar-cane, 373 sqq.
— for tea, 298, 300 sqq.
— on barley in S. Australia, 327–32.
— on wheat in S. Australia, 193–205, 325–7.
Australia, red soils in, 106.
— soil surveys in, 333–40.
Australia, South, climate of, 193–5.
— manuring of wheat and barley in, 193–206, 325–32.
Australia, Queensland, manuring of sugar-cane in, 378–9.

Bacon, possible production of in Great Britain, 216–17.
Barbados, red soils in, 105.
Barley, S. Australia, effect of N-fertilizers on, 327–32.
— yield of, after legumes, 27 sqq.
Basic slag, for sugar-cane, 375.
— for tea, 298.
Beans, residual manurial value of, 22 sqq.
Bermuda, lily industry in, 213.
Blackman, G. E., botanical composition of intensively managed pastures, 253–70.
Butter, carotene, xanthophyll, and vitamin contents of, 71, 74, 81.
— colour of and vitamins in, 68 sqq.

Canada, breeding of early-ripening spring wheat in, 3–16.
Canada, Western, soil surveys in, 33–42.
Carotene, in grass and butter, 68 sqq.
Cashmore, W. H., haymaking machinery, 58–67.
Cereals, experiments with in N. Nigeria, 113–21.
— growing of, in Kent in 13th century, 82–4.
Ceylon, imports of N-fertilizers, 299.
— tea-cultivation in, 245 sqq., 297 sqq.
Chittenden, R. J., asexual propagation of *Hevea*, 307–15.
Clovers, residual manurial value of, 22.
Coffea arabica root-systems, 270–84, 285–96; effect of cultivation on, 291–3; of high water-table on, 288; of manuring on, 291; of overbearing on, 293–5; of soil conditions on, 286 sqq.; of soil reaction on, 289–91.
— isolation of, 271–4.
— typical, 285–6.
Cows, feeding-experiments with, 69.
Cuba, manuring of sugar-cane in, 377.
— red soils in, 105.
Cyprus, climate of, 87, 96, 156.
— soil profiles from, 85–96, 156–64.
— soil map of, 86.
— soils of, 160–4.
— weathering of igneous rocks in, 92–5, 156–60; of sedimentary rocks in, 88–92.

Diabase, weathering of, in Cyprus, 156; in French Guinea, 164; in Spitzbergen, 164.
Diammonium phosphate, on wheat in S. Australia, 201–2.
Disease, aster yellows, 208, 209.
— curly top of sugar-beet, 208, 209, 212.
— dieback of coffee, 275, 293–5.
— *Haemonchus contortus* in lambs, 17–21.
— *Krauselkrankheit* in Germany, 207, 213.
— leaf-curl of cotton in Sudan, 206, 213.
— mosaic, 207 sqq.
— mosaic of sugar-cane, 212.
— mosaic of tobacco, 207, 211.
— potato virus X, 208.
— rosette of ground-nuts, 213.
— rosette of wheat in N. America, 207.
— virus in plants, 206–14.
— yellows of peach-tree, 208.
Dodds, H. H., manuring of sugar-cane, 368–80.
Drummond, J. C., *see* Watson, S. J.

Eden, T., manuring of tea, 297–306.
Elliot, Rt. Hon. W. E., 2.
Ellis, J. C. B., *see* Watson, J. A. S.
Evans, I. B. Pole, agricultural possibilities of some African grasses, 341–6.
Ewbank, Elinor K., *see* Reifenberg, A.

Farm-yard manure, experiments with, in N. Nigeria, 113–21.
Fertilizers for sugar-cane, 372–5.
— effects on wheat, 193–205, 325–7.
— for tea, 298 sqq.
— nitrogen and phosphate on pastures, 51, 70, 122, 253, 361, 364.
Field experiments, analysis of replicated, 129–42.

Filter-cake, as manure for sugar-cane, 369–70.
Fraser, A. H. H., nutritional state of sheep and susceptibility to stomach worm, 17–21.

Gabbro and gabbro-norite, weathering of, in Cyprus, 156, 164.
Grass, artificial drying of, 68.
— artificially dried, as food for cows, 63; composition of, 70.
— Elephant, in S. and E. Africa, 345.
— Kikuyu, in S. Africa, 345.
— silage, 70.
Grasses, agricultural possibilities of some African, 341–6.
— *Digitaria*, in S. Africa, 342–5.
— pasture, experiments with, 49–56, 122–7, 224–8, 253–70, 361–7; seasonal productivity curves of, 265.
Grassland, management of, 43–57, 122–8, 223–34, 253–70, 361–7.
— herbage: creeping plants in, 230–1.
— — dry-matter in, 358.
— — edaphic and biotic influences on, 366–7.
— — effect of autumn protection on, 228–9; of 'burn' on palatability of, 352, 356–7; of cutting on roots of, 46; of defoliation on, 45; of grazing on composition of, 268; of mowing on spring growth of, 48; of over- and undergrazing on, 125; of winter grazing on, 48.
— — in poor pastures, 361–6.
— — in winter, effect of summer and autumn grazing on, 47.
— — on Welsh open-hill grazings, 353–5.
— — palatability of grasses and clovers in lowland, 350–2; of miscellaneous herbs in, 347–60.
— — start of active growth of, 49.
— rotational grazing of, 124, 253 sqq., 363; effect on botanical composition of, 253–70.
— weed-control in, 56.
Green manuring of tea, 299.
Greenwood, M., *see* Hartley, K. T.
Guinea corn, experiments with, N. in Nigeria, 113–21.
Gurney, H. C., *see* Richardson, A. E. V.

Hard-pan in E. African soils, 276, 286–7.
Hardy, F., cultivation properties of tropical red soils, 103–12.
Hartley, K. T., effect of small applications of f.y.m. on cereals in Nigeria, 113–21.
Hawaii, manuring of sugar-cane in, 376.
— red soils in, 103.
Haymaking, machines for, 58–67.
Heilbron, I. M., *see* Watson, S. J.
Herbs, miscellaneous, in corn-stubbles, 349–50; in lowland grassland, 348.
Hevea, asexual propagation of, 307–15.
Hevea, budding technique, 312–14.
— lay-out of bud-grafted clearings, 309–10.
— nurseries and multiplication-beds of bud-wood, 311.
— proving of clones, 308–9.
— selection of bud-parents, 307; of clones, 309.
— tapping systems of bud-grafted, 310–11.

India, manuring of sugar-cane in, 378.
— red soils in, 107.
— tea-cultivation in, 245 sqq., 297 sqq.

Java, manuring of sugar-cane in, 375; of tea in, 297 sqq.
Joel, A. H., soil surveys in W. Canada, 33–42.
Jones, M. G., grassland management, 43–57, 122–8, 223–4, 361–7.

Keen, B. A., experimental study of soil cultivation, 97–102.

Latin squares, formation of, 235–44.
Legumes, residual manurial value of, 22–32.
Lime in tea soils, effect of, 248.
Limestone, weathering of, 160–3.
Lucerne, residual manurial value of, 22 sqq.

Machines, for haymaking, 4, 58–67.
— tractor, 60–3.
Mann, H. H., climatic and soil requirements of tea, 245–52.
Mauritius, manuring of sugar-cane in, 378–9.
Milk, from cows fed on A.I.V.-fodder, 151.
— from cows fed on dried grass and grass silage, yield and composition of, 76–8.
Miller, William C., inheritance of wool-characters in sheep, 173–92.
Milton, W. E. J., palatability of the self-establishing species contributing to different types of grassland, 347–60.
Molasses, as manure for sugar-cane, 370–2.
Morton, R. A., *see* Watson, S. J.
Murray, K. A. H., pig industry in Great Britain, 215–22.

Newman, J. E., *see* Cashmore, W. H.
Newman, L. H., early-ripening wheats in Canada, 3–16.
Nicol, Hugh, residual values of leguminous crops, 22–32.
Nitro-chalk on grassland, 49, 122, 253, 364.
— on wheat in S. Australia, 201–2.
Nutman, F. J., root-system of *Coffea arabica*, 270–84, 285–96.

Oats, yield of, after legumes, 27 sqq.
Organic manures, compost for yams, in Trinidad, 324.
— f.y.m. in N. Nigeria, 113.

Organic manures, oilseed cakes, &c., for tea, 298.
— pen manure, 368–9.

Peas, residual manurial value of, 22 sqq., 325–7.
Pelham, R. A., soils and grain-growing in Kent in 13th century, 82.
pH, influence of, on protein-content of silage, 144.
Philippine Islands, manuring of sugar-cane in, 379.
Pig industry in Great Britain, future of, 214–22; pig numbers in Great Britain, 216, 218–22; prices and feeding-stuffs, ratio of, 218–22.
Potash for tea, 301, 302, 303.
— for sugar-cane, 370–2, 374, 376, 377, 378, 379.
Prescott, J. A., soil surveys in Australia, 333–40.
Puerto Rico, manuring of sugar-cane in, 376.

Reifenberg, A., soil profiles from Cyprus, 85–96, 156–64.
Richardson, A. E. V., nitrogenous fertilizers on wheat and barley in S. Australia, 193–205, 325–32.
Robertson, David, *see* Fraser, A. H. H.

Sainfoin, residual manurial value of, 22 sqq.
Season, effect of, on residual manurial values of legumes, 31.
Serpentine, weathering of, in Cuba, 163; in Cyprus, 92–5, 163; in Germany, 163.
Sheep, food-consumption of fattening, 165–72.
— inheritance of wool-characters in, 173–92.
— nutritional state and susceptibility to stomach worm, 17–21.
Silage, A.I.V.-method, 143–55.
— grass, composition of, 70.
Skilbeck, D., *see* Watson, J. A. S.
Smith, J. Henderson, virus diseases in plants, 206–14.
Sodium nitrate for sugar-cane, 374; for tea, 302; on wheat in S. Australia, 201–2.
Soil, at Samaru, N. Nigeria, analysis of, 114.
— cultivation, experimental study of, 97.
— formation, in Cyprus, 85, 156, 160.
— nitrate-content of, fallowed and cropped, in S. Australia, 194 sqq.; in Nigeria, 114 sqq.
— survey, in W. Canada, methods of, 35–40; in Australia, methods of, 336–8.
Soils, in Australia, classification of, 338–40; survey of, 333–40.
— in W. Canada, classification of, 40; survey of, 33–42.
— coffee, in British E. Africa, 275–84, 286–8; reaction of, 274, 289–91.
Soils, hard-pan in British E. Africa, 286–7.
— in Cyprus, 160–4.
— lateritic red, composition of, 110–11.
— red, in Cyprus, 85, 156, 160; in E. Africa, 276.
— tropical red, 103–12; classification of, 107–8; from sedimentary rocks, 108, 109.
— tea, requirements of, 245–52.
Stomach worm (*Haemonchus contortus*) in sheep, 17–21.
Sugar-cane, causes of increased production of, 368–9.
— manurial practices, 375–9.
— manuring of, 368–80; with artificial manures, 372–5; with filter-cake, 369–70; with molasses, 370–2; with pen manure, 368–9.
Superphosphate for sugar-cane, 375; for tea, 298, 300 sqq.
— on barley in S. Australia, 327, 332.
— on wheat in S. Australia, 201–2, 325–7.

Tea, climatic and soil requirements of, 245–52.
— effect of manuring on quality of, 249–51, 304–5.
— manuring of, 297–306.
— yield-increases from N-manures, 302.
Tea-cultivation, geographical distribution of, 245–6.
Tea soils, acidity of, 248–9.
— plant-foods required by, 249–51.
Terra rossa, 88 sqq., 109, 160.
Thistle, creeping, control of, in pastures, 56.
Thomas, Rt. Hon. J. H., 1.
Trinidad, experiments with yams in, 316–24.

Urea, on wheat in S. Australia, 201–2.

Vetch, residual manurial value of, 22.
Virtanen, A.I., A.I.V.-method of preserving fresh fodder, 143–55.
Virus diseases in plants, 206–14; classification of, 209–10; control of, 211–13; insect carriers of, 208; resistance to, 211–13; transmission of, 207–8, 213.
Virus organisms, nature of, 210–11; size of, 211–12; specificity of, 208–9.
Vitamins in grass and butter, 68 sqq.

Watson, J. A. S., food-consumption of fattening sheep, 165–72.
Watson, S. J., influence of dried grass in winter ration of dairy cow on colour of, and vitamins A and D in butter, 68–81.
West Indies, British, manuring of sugar-cane in, 377–8.
Wheat, early-ripening varieties in Canada, 3–16.

Wheat, S. Australia, effect of N-fertilizers on, 193–205, 325–7; of previous crop in response to N, 325–7.
— inheritance of earliness in, 13.
— yield of, after legumes, 25, 32.
Wood, R. Cecil, experiments with yams in Trinidad, 316–24.
Wool, sheep's, desiderata for genetic experiments on, 173.
— — inheritance of characters of, 173–92; of crimp, 181; of fibre-length of, 179; of fineness of, 175; of kemp fibres of, 183; of weight and density of fleece, 182.
Xanthophyll, in butter, 68, 74, 81.

Yams, in Trinidad, experiments with, 316–24; early *v.* late planting, 322; cultural experiments, 319–24; increased yields from closer planting, 319; manurial experiments, 320–1; planting time, 323–4; use of organic compost, 324; varietal experiments, 316–19.
Yates, F., analysis of replicated field experiments when results are incomplete, 129–42.
— formation of Latin squares for field experiments, 235–44.

PRINTED IN GREAT BRITAIN AT THE UNIVERSITY PRESS, OXFORD
BY JOHN JOHNSON, PRINTER TO THE UNIVERSITY